

EXCELLENCE IN IMMEDIACY

Straumann® Bone Level Tapered Implant
Unlock the magic behind
immediate placement.



In combination with:



“I have found the BLT design extremely beneficial for my single and multiple tooth immediate provisional cases. The new tapered apical design allows for very predictable insertion torques of 35 Ncm and excellent primary stability.”

Brad McAllister, DDS, PhD
Periodontal Associates, Tualatin, Oregon, USA



ROXOLID®
Stronger than titanium.¹



SLACTIVE®
Superb healing capabilities.^{2, 3, 4, 5}

Legendary versatility and predictability even in very challenging situations.

As global market needs for dental implants are shifting towards shorter treatment times, clinicians expect solutions that reduce chair time, improve efficiency, reliability and profitability, while meeting the needs of specific clinical and anatomic situations. At the same time, patients are demanding solutions that deliver esthetic results faster.

Trust the powerful combination of the Straumann® Bone Level Tapered Implant design, with the high-performance surface, SLActive® and the high-strength material, Roxolid® designed to maximize primary and secondary stability in all bone classes making them ideal for immediate placement and loading* even in very challenging protocols.



Extraction sockets



Anatomical limitations

* When good primary stability is achieved.



APICALLY TAPERED

Designed for enhanced
primary stability.⁶



IMMEDIATE TREATMENT

Designed for immediate
placement and loading.⁶



GUARANTEE

Roxolid® Lifetime Plus Guarantee.
Our quality, your assurance.

Straumann® Bone Level Tapered Implant

Digital workflows

Designed for immediate restoration

- Flexible milling options
- Full digital connectivity



| Roxolid®

Reducing invasiveness with smaller implants

- Preserves vital structures and vascularization^{7, 8}
- Reduces invasive grafting procedures^{7, 8}
- Increases treatment options in challenging anatomical situations and narrow interdental spaces^{8, 9}



| SLActive®

High performance surface designed to:

- Reduce healing time to 3–4 weeks^{2, 3}
- Deliver high long-term predictability in immediate loading¹⁰
- Provide outstanding success rates in compromised patients^{4, 5, 11, 12}
- Promote bone regeneration, even at compromised sites^{13, 14, 15, 16}



| Apically tapered

Unrivalled primary stability even in compromised bone situations

- Full-depth thread to apex for early engagement
- Self-cutting in underprepared sites
- Protecting anatomical structure with round tip





Prosthetic components

- Single- and multi-unit replacements: screw- or cement-retained
- Edentulous treatment: fixed or removable options
- CrossFit® Connection: prosthetic components and simplified handling from the Bone Level System
- Consistent emergence profiles™: the bone level standard for enhanced bone preservation and esthetics



Bone Control Design™

- Maximized crestal bone preservation¹⁷
- Respects the biological distance/width and microgap control¹⁸
- Optimal position of smooth and rough surface interface^{15, 18}
- Biomechanical implant design and implant surface osteoconductivity¹⁹



Crossfit® connection

Combining easy handling and high reliability

- Self-guiding connection and four internal grooves provide easy handling and deliver confidence for component positioning
- Exact implant-abutment fit and optimized stress distribution ensure long-mechanical stability
- Internal cone allows for greater flexibility in prosthetic treatments

Reaching new dimensions. Small, strong and enduring.

The SmallOne Bone Level Tapered Implant 2.9 mm is specifically engineered to address esthetic challenges in compromised situations such as narrow interdental spaces, or reduced bone availability. Straumann BLT 2.9 mm implants allow you to treat your patients with confidence and give them a less invasive solution with predictable outcomes.



LESS INVASIVE

The perfect combination of shape, strength and healing power.⁶



NATURAL-LOOKING ESTHETICS

Novel prosthetic shape designed to match the anatomy of natural teeth.



FULL STRENGTH

A strong and truly reliable treatment option thanks to Roxolid®.¹



REFERENCES

- 1 Medvedev AE, Molotnikov A, Lapovok R, Zeller R, Berner S, Habersetzer P, Dalla Torre F. Microstructure and mechanical properties of Ti-15Zr alloy used as dental implant material. *J Mech Behav Biomed Mater*. 2016 Sep;62:384-398. doi: 10.1016/j.jmbbm.2016.05.008. Epub 2016 May 11. **2** Lang NP, Salvi GE, Huynh-Ba G, Ivanovski S, Donos N, Bosshardt DD. Early osseointegration to hydrophilic and hydrophobic implant surfaces in humans. *Clin Oral Implants Res*. 2011 Apr;22(4):349-56. doi:10.1111/j.1600-0501.2011.02172.x **3** Oates TW, Valderrama P, Bischof M, Nedir R, Jones A, Simpson J, Toutenburg H, Cochran DL. Enhanced implant stability with a chemically modified SLA surface: a randomized pilot study. *Int J Oral Maxillofac Implants*. 2007 Sep-Oct;22(5):755-60 **4** Nelson, K., Stricker, A., Raguse, J.-D. Nahles, S. (2016), Rehabilitation of irradiated patients with chemically modified and conventional SLA implants: a clinical clarification. *J Oral Rehabil*, 43: 871–872. doi:10.1111/joor.12434. **5** Heberer S, Kilic S, Hossamo J, Raguse J-D, Nelson K. Rehabilitation of irradiated patients with modified and conventional sandblasted, acid-etched implants: preliminary results of a split-mouth study. *Clin. Oral Impl. Res*. 22, 2011; 546–551. **6** Dard M, Kuehne S, Obrecht M, Grandin M, Helfenstein J, Pippenger BE. Integrative Performance Analysis of a Novel Bone Level Tapered Implant. *Adv Dent Res*. 2016 Mar;28(1):28-33. doi: 10.1177/0022034515624443. **7** Ioannidis A, Gallucci GO, Jung RE, Borzangy S, Hammerle CH, Benic GI. Titanium-zirconium narrow-diameter versus titanium regular-diameter implants for anterior and premolar single crowns: 3-year results of a randomized controlled clinical study. *J Clin Periodontol*. 2015 Nov;42(11):1060-70. doi: 10.1111/jcpe.12468. Epub 2015 Nov 14. **8** Al-Nawas B, Domagala P, Fragola G, Freiburger P, Ortiz-Vigón A, Rousseau P, Tondela J. A Prospective Noninterventional Study to Evaluate Survival and Success of Reduced Diameter Implants Made From Titanium-Zirconium Alloy. *J Oral Implantol*. 2015 Aug;41(4):e118-25. doi: 10.1563/AAID-JOI-D-13-00149. Epub 2014 Mar 25. **9** Altuna P, Lucas-Taulé E, Gargallo-Albiol J, Figueras-Álvarez O, Hernández-Alfaro F, Nart J. Clinical evidence on titanium-zirconium dental implants: a systematic review and meta-analysis. *Int J Oral Maxillofac Surg*. 2016 Jul;45(7):842-50. doi: 10.1016/j.ijom.2016.01.004. Epub 2016 Feb 3. **10** Nicolau P, Guerra F, Reis R, Krafft T, Benz K, Jackowski J 10-year results from a randomized controlled multicenter study with immediately and early loaded SLActive implants in posterior jaws. Presented at 25th Annual Scientific Meeting of the European Association of Osseointegration – 29 Sep – 1 Oct 2016, Paris. **11** Cabrera-Domínguez J, Castellanos-Cosano L, Torres-Lagares D, Machuca-Portillo G. A Prospective Case-Control Clinical Study of Titanium-Zirconium Alloy Implants with a Hydrophilic Surface in Patients with Type 2 Diabetes Mellitus. *Int J Oral Maxillofac Implants*. 2017 Sep-Oct;32(5):1135-1144. doi: 10.11607/jomi.5577; **12** Cabrera-Domínguez J. A prospective, two-year clinical trial of titanium-zirconium alloy implants (Roxolid® Straumann®) with hydrophilic surface (SLActive®) in patients with Type 2 Diabetes Mellitus. presented during 26th Annual Scientific Meeting of the European Association of Osseointegration – 5-7 Oct 2017, Madrid, Spain **13** Kuchler U, Chappuis V, Bornstein mm, Siewczyk M, Gruber R, Maestre L, Buser D. Development of Implant Stability Quotient values of implants placed with simultaneous sinus floor elevation – results of a prospective study with 109 implants. *Clin. Oral Impl. Res*. 28, 2017, 109–115 **14** Schwarz, F., et al., Bone regeneration in dehiscence-type defects at chemically modified (SLActive®) and conventional SLA® titanium implants: a pilot study in dogs. *J Clin.Periodontol*. 34.1 (2007): 78–86 **15** Schwarz, F., et al., Bone regeneration in dehiscence-type defects at non-submerged and submerged chemically modified (SLActive®) and conventional SLA titanium implants: an immunohistochemical study in dogs. *J Clin.Periodontol*. 35.1 (2008): 64–75. **16** Straumann (2016). SLActive supports enhanced bone formation in a minipig surgical GBR model with coronal circumferential defects. Unpublished data. **17** French D, Cochran DL, Ofec R. Retrospective Cohort Study of 4,591 Straumann Implants Placed in 2,060 Patients in Private Practice with up to 10-Year Follow-up: The Relationship Between Crestal Bone Level and Soft Tissue Condition. *Int J Oral Maxillofac Implants*. 2016 Nov/Dec;31(6):e168-e178. doi: 10.11607/jomi.4932. **18** Cochran DL, Mau LP, Higginbottom FL, Wilson TG, Bosshardt DD, Schoolfield J, Jones AA. Soft and hard tissue histologic dimensions around dental implants in the canine restored with smaller-diameter abutments: a paradigm shift in peri-implant biology. *Int J Oral Maxillofac Implants*. 2013 Mar-Apr;28(2):494-502. doi: 10.11607/jomi.3081. **19** Cochran D, Stavropoulos A, Obrecht M, Pippenger B, Dard M. A Comparison of Tapered and Nontapered Implants in the Minipig. *Int J Oral Maxillofac Implants*. 2016 Nov/Dec;31(6):1341-1347. doi: 10.11607/jomi.4712.



International Headquarters

Institut Straumann AG

Peter Merian-Weg 12

CH-4002 Basel, Switzerland

Phone +41 (0)61 965 11 11

Fax +41 (0)61 965 11 01

www.straumann.com

© Institut Straumann AG, 2018. All rights reserved.

Straumann® and/or other trademarks and logos from Straumann® mentioned herein are the trademarks or registered trademarks of Straumann Holding AG and/or its affiliates.

 **straumann**