

Straumann® Emdogain®

Wellbeing and
faster recovery after
oral surgery

Straumann®
Emdogain®
More than 2 million
patients treated



Esthetics, comfort and efficiency: Straumann® Emdogain®

The first days after oral surgery – such as implant placement, bone augmentation, tooth extraction, gum surgery or periodontal surgery – bear the risk of pain, swelling and discomfort which may affect your quality of life. Moreover, wound healing complications may occur during this period which may have an impact on the esthetics of the operated area.



LESS PAIN AND SWELLING AFTER SURGERY⁴

Emdogain® reduces the incidence of pain and swelling following oral surgical procedures*

- **5 times** more patients report no pain after one week
- **7 times** more patients report no swelling after one week

*compared to treatment with membranes

Straumann® Emdogain® is a unique gel which contains specific natural proteins. When applied during surgery, Straumann® Emdogain® stimulates cells and biological processes that are fundamental for wound healing¹ and thereby

- Accelerates wound closure²
- Reduces the risk for post surgical pain and swelling^{3,4}
- Reduces the risk of wound complications by stimulating the healing of gum tissue^{2,5} and protecting against oral pathogens⁶
- Improves the esthetic result by accelerating gum tissue formation and maturation^{7,8}

WHY ENDURE PAIN?

You deserve wellbeing and fast recovery. Ask your dentist for more information on Straumann® Emdogain®.

REFERENCES

- 1 Miron RJ, Dard M, Weinreb M. Enamel matrix derivative, inflammation and soft tissue wound healing. *J Periodontol Res.* 2015 Oct;50(5):555-69. doi: 10.1111/jre.12245. 2 Villa O, Wohlfahrt JC, Mdlá I, Petzold C, Reseland JE, Snead ML, Lyngstadaas SP. Proline-Rich Peptide Mimics Effects of Enamel Matrix Derivative on Rat Oral Mucosa Incisional Wound Healing. *J Periodontol.* 2015 Dec;86(12):1386-95. doi: 10.1902/jop.2015.150207. 3 Ozcelik O, Haytac MC, Seydaoglu G. Immediate post-operative effects of different periodontal treatment modalities on oral health-related quality of life: a randomized clinical trial. *J Clin Periodontol.* 2007 Sep;34(9):788-96. 4 Jepsen S, Heinz B, Jepsen K, Arjomand M, Hoffmann T, Richter S, Reich E, Sculean A, Gonzales JR, Bödeker RH, Meyle J. A randomized clinical trial comparing enamel matrix derivative and membrane treatment of buccal Class II furcation involvement in mandibular molars. Part I: Study design and results for primary outcomes. *J Periodontol.* 2004 Aug;75(8):1150-60. 5 Guimarães GF, de Araújo VC, Nery JC, Peruzzo DC, Soares AB. Microvessel Density Evaluation of the Effect of Enamel Matrix Derivative on Soft Tissue After Implant Placement: A Preliminary Study. *Int J Periodontics Restorative Dent.* 2015 Sep-Oct;35(5):733-8. 6 Arweiler NB, Auschill TM, Donos N, Sculean A. Antibacterial effect of an enamel matrix protein derivative on in vivo dental biofilm vitality. *Clin Oral Investig.* 2002 Dec;6(4):205-9. 7 Piloni A, Paolantonio M, Camargo PM. Root coverage with a coronally positioned flap used in combination with enamel matrix derivative: 18-month clinical evaluation. *J Periodontol.* 2006 Dec;77(12):2031-9. 8 Tonetti MS, Fourmoussis I, Suvan J, Cortellini P, Brägger U, Lang NP, European Research Group on Periodontology (ERGOPERIO). Healing, post-operative morbidity and patient perception of outcomes following regenerative therapy of deep intrabony defects. *J Clin Periodontol.* 2004 Dec;31(12):1092-8.

International Headquarters

Institut Straumann AG

Peter Merian-Weg 12

CH-4002 Basel, Switzerland

Phone +41 (0)61 965 11 11

Fax +41 (0)61 965 11 01

www.straumann.com

© Institut Straumann AG, 2017. All rights reserved.

Straumann® and/or other trademarks and logos from Straumann® mentioned herein are the trademarks or registered trademarks of Straumann Holding AG and/or its affiliates.